

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
ONR-TR-26	AD-AD99	3. RECIPIENT'S CATALOG NUMBER  992	
Organometallic Photochemist  Interface. Photochemistry o	f Surface-Confined Cob-	5. TYPE OF REPORT & PERIOD COVERE  Interim Technical Report	
alt Tetracarbonyl Monitored Infrared Photoacoustic Spec		5. PERFORMING ORG. REPORT NUMBER	
John B. Kinney, Ralph H. St and Mark S. Wrighton	aley, Carol L. Reichel,	NOO14-75-C-0880	
PERFORMING ORGANIZATION NAME AND Department of Chemistry, Rm Massachusetts Institute of Cambridge, Massachusetts 0	. 6-335 Technology	19. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 051-579	
11. CONTROLLING OFFICE NAME AND ADDR	RESS .	12. REPORT DATE	
Office of Naval Research Department of the Navy Arlington, Virginia 22217		May 11, 1981 13. NUMBER OF PAGES 20	
14. MONITORING AGENCY NAME & ADDRESS	(If different from Controlling Office)	18. SECURITY CLASS. (of this report)	
		UNCLASSIFIED	
,		15a. DECLASSIFICATION/DOWNGRADING	

Approved for public release; reproduction is permitted for any purpose of the United States Government; distribution unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Distribution of this document is unlimited.

18. SUPPLEMENTARY NOTES

Prepared and accepted for publication in the Journal of the American Chemical Society.

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Photoacoustic spectroscopy, organometallic photochemistry, surface photochemistry, interfacial photochemistry

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

(Please turn over for Abstract.)

DD 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE

UNCLASS I FIED

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

High surface area  $SiO_2$  (400 m<sup>2</sup>/g) functionalized first with  $\longrightarrow$ SiH by reaction with (EtO) 3SiH and then reacted with Co2(CO) 8 yields SiCo(CO)4  $(S = SiO_2)$ . Photochemistry of  $S = SiCo(CO)_4$  exposed to various gases has been studied by using Fourier transform infrared photoacoustic spectroscopy, FTIR/PAS. Near-uv irradiation under  $^{13}$ CO yields  $\text{S} = \text{SiCo(CO)}_{n} (^{13}\text{CO)}_{d-n}$  and the reaction can be reversed by irradiation under CO. Irradiation under  $0_2$ yields loss of metal carbonyl signals and  ${\rm CO}$  and  ${\rm CO}_2$  are observed as gas phase products. Experiments with  $^{18}O_2$  and with  $^{13}CO$ -enriched  $\bigcirc$  SiCo(CO)<sub>4</sub> show that  $CO_2$  originates from CO initially bound to CO and  $O_2$  in the gas phase. Ligand substitution by gas phase PF3, ethylene, propylene, isobutylene, or propyne can be detected when  $S = SiCo(CO)_4$  is irradiated in the presence of the gas. All photochemistry is consistent with loss of CO as the primary chemical result of photoexcitation of  $S = SiCo(CO)_4$ . The FTIR/PAS technique is useful on sample sizes of ~1 mg and gives infrared spectra of comparable quality to transmission infrared of KBr/ (S)-SiCo(CO)<sub>4</sub> pellets. The FTIR/PAS technique requires no pretreatment or manipulation of the sample and the monitoring of surface and gas phase products can be done in situ.

OFFICE OF NAVAL RESEARCH

12)22

CONTRACT NO0014-75-C-0880

/ミノ

Task No. NR 051-579

TECHNICAL REPORT, NO. 26

/ )

ORGANOMETALLIC PHOTOCHEMISTRY AT THE SOLID/GAS INTERFACE. PHOTOCHEMISTRY

OF SURFACE-CONFINED COBALT TETRACARBONYL MONITORED BY FOURIER TRANSFORM

INFRARED PHOTOACOUSTIC SPECTROSCOPY

Ьy

John B./Kinney, Ralph H./Staley, Carol L./Reichel, and Mark S./Wrighton

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Prepared for publication in the Journal of the American Chemical Society

May 11, 1981

11 77 11987

Reproduction in whole or in part is permitted for any purpose of the United States Government

This document has been approved for public release and sale; its distribution is unlimited.

· 1011/11/11

Organometallic Photochemistry at the Solid/Gas Interface. Photochemistry
of Surface-Confined Cobalt Tetracarbonyl Monitored by Fourier Transform
Infrared Photoacoustic Spectroscopy

Abstract: High surface area SiO<sub>2</sub> (400 m<sup>2</sup>/g) functionalized first with SiH by reaction with  $(Et0)_3$ SiH and then reacted with  $Co_2(C0)_8$  yields  $S = SiCo(CO)_4$  ( $S = SiO_2$ ). Photochemistry of  $S = SiCo(CO)_4$  exposed to various gases has been studied by using Fourier transform infrared photoacoustic spectroscopy, FTIR/PAS. Near-uv irradiation under 13CO yields  $S \rightarrow SiCo(CO)_n(^{13}CO)_{4-n}$  and the reaction can be reversed by irradiation under CO. Irradiation under O2 yields loss of metal carbonyl signals and CO and  $CO_2$  are observed as gas phase products. Experiments with  $^{18}O_2$  and with  $^{13}$ CO-enriched (S)=SiCo(CO)<sub>4</sub> show that CO<sub>2</sub> originates from CO initially bound to Co and  $0_2$  in the gas phase. Ligand substitution by gas phase PF<sub>3</sub>, ethylene, propylene, isobutylene, or propyne can be detected when  $S) > SiCo(CO)_4$  is irradiated in the presence of the gas. All photochemistry is consistent with loss of CO as the primary chemical result of photoexcitation of (S) SiCo(CO)<sub>4</sub>. The FTIR/PAS technique is useful on sample sizes of ~1 mg and gives infrared spectra of comparable quality to transmission infrared of KBr/(S)-SiCo(CO)<sub>4</sub> pellets. The FTIR/PAS technique requires no pretreatment or manipulation of the sample and the monitoring of surface and gas phase products can be done in situ.

Organometallic Photochemistry at the Solid/Gas Interface. Photochemistry
of Surface-Confined Cobalt Tetracarbonyl Monitored by Fourier Transform
Infrared Photoacoustic Spectroscopy

John B. Kinney, Ralph H. Staley, Carol L. Reichel, and Mark S. Wrighton\*

Department of Chemistry

Massachusetts Institute of Technology

Cambridge, Massachusetts 02139 U.S.A.

we wish to report the photochemistry of surface-confined -Co(CO)<sub>4</sub> fragments exposed to reactive gases. Such photochemistry is of importance in establishing primary events following excitation of the molecular entity attached to the surface and is of relevance to the photoactivation of surface-confined catalysts. We apply, for the first time, the technique of Fourier transform infrared photoacoustic spectroscopy (FTIR/PAS)<sup>3</sup> to monitor the photoreactions of a species on the surface including in situ monitoring of reactions involving a gas phase species. Without any sample manipulation or preparation, this technique has allowed characterization of the photochemistry of surface species with the molecular level specificity generally possible when infrared absorption spectroscopy is used to monitor reactions of metal carbonyls in homogeneous solution. The results herein establish FTIR/PAS as a technique of unequalled capability in monitoring such surface chemistry.

The system studied is  $S \to SiCo(CO)_4$  where  $S \to Tepresents$  a high surface area silica (~400 m²/g from Alfa). The synthetic procedure follows from known reaction chemistry and was carried out according to the representation in (1) and (2). The functionalization of metal oxides such

surface-OH 
$$\frac{25^{\circ}C}{(Et0)_{3}SiH}$$
 surface-O-SiH (1)

as  $\mathrm{SiO}_2$  using  $(\mathrm{RO})_3\mathrm{SiR}^1$  reagents and reaction of  $\mathrm{Co}_2(\mathrm{CO})_8$  with  $\mathrm{R}_3\mathrm{SiH}$  to give  $\mathrm{R}_3\mathrm{SiCo}(\mathrm{CO})_4$  are both well-known. The reaction to prepare SiH is established by transmission infrared of KBr/SiH (10/1 by weight) pellets showing a feature at ~2250 cm<sup>-1</sup> associated with the Si-H stretch; no feature in this region is observed for the  $\mathrm{SiO}_2$  prior to treatment with (EtO)\_3SiH. Absorption is also observed in the C-H region reflecting the fact that some residual EtO— groups remain. Reaction of SiH with  $\mathrm{Co}_2(\mathrm{CO})_8$  results in additional infrared absorptions in the metal carbonyl region at ~2110 (rel. o.d. = 0.3); 2050 (rel. o.d. = 0.5); and ~2020 (rel. o.d. = 1.0) cm<sup>-1</sup>. The number of bands and their relative intensities accord well with bands expected for SiCo(CO)\_4 based on species such as (EtO)\_3SiCo(CO)\_4. From a combination of infrared and elemental analyses the coverage of  $-\mathrm{SiCo}(\mathrm{CO})_4$  is sub-monolayer (~10<sup>-11</sup> mol/cm<sup>2</sup>), and there is generally considerable  $-\mathrm{SiH}$  remaining after reaction with  $\mathrm{Co}_2(\mathrm{CO})_8$ .

Figure 1 includes a typical FTIR/PAS characterization of ~1 mg of  $S-SiH/-SiCo(CO)_4$ . The relative FTIR/PAS signals are proportional to infrared absorbances found for  $S-SiCo(CO)_4$  in KBr. For both forms of infrared the metal carbonyl signals are broader and less well-resolved than the spectral features for species such as  $R_3SiCo(CO)_4$  in alkane solution. The FTIR/PAS of the  $S-SiH/-SiCo(CO)_4$  is just as good in terms of signal-to-noise as the FTIR/absorption and does not require palleting or other pretreatment of the sample. Quantities of sample required for FTIR/PAS are as small, or smaller, as typically used to prepare 1 cm diameter KBr  $S-SiH/-SiCo(CO)_4$  pellets.

The unique capabilities of the FTIR/PAS are revealed in the photo-chemical studies. Typically, a ~1 mg sample of  $S - SiH/-SiCo(CO)_4$  is placed in the FTIR/PAS cell and irradiated at 25°C with the 300-400 nm output from a 150 W Xe lamp focused to ~1 cm<sup>2</sup>; the input optical power to the sample

is-100 mW/cm<sup>2</sup>. The undisturbed sample can then be characterized by FTIR/PAS.<sup>8</sup> The first significant experiment concerns the results from irradiation of  $S=SiH/=SiCo(CO)_4$  when the cell contains 1 atm of  $SiH/=SiCo(CO)_4$  when the cell contains 1 atm of SiH/

$$s = sico(co)_4 + {}^{13}co \xrightarrow{hv} s = sico(co)_n ({}^{13}co)_{4-n}$$
 (3)

$$S = SiCo(CO)_4 \xrightarrow{hv} S = SiCo(CO)_3 + CO$$
 (4)

excitation. When irradiation is carried out for short periods of time, spectral changes are consistent with the sequential replacement of bound CO by  $^{13}$ CO. The FTIR/PAS monitoring can be carried out without pumping out the cell, using the reactive gas as the acoustic coupler. When the gases are not pumped out the  $^{13}$ CO is detected and  $^{12}$ CO is detected as a gas phase photoproduct, as expected. Additionally, small amounts of  $CO_2$  and  $CO_2$ are detected as photoproducts (cf. below). Irradiation of (S)  $Si(CO)_n(^{13}CO)_{4-n}$ for  $n \approx 0$  under 1 atm CO results in the regeneration of with ~80% recovery of the FTIR/PAS signal for the carbonyl species. Chemically, these results establish CO loss as the principal result of photoexcitation of the (S)-SiCo $(CO)_4$ . This finding parallels the conclusion drawn from studies of  $R_3SiCo(CO)_4$  in homogeneous solution and  $(S)_4$ SiCo(CO)\_4 suspended in solution. In particular, the FTIR/PAS shows little or no importance for photoinduced cleavage of the Si-Co bond that would lead to  $Co(CO)_4$  or to the dimer  $Co_2(CO)_8$ . With the FTIR/PAS technique, the data are much more easily obtained and without sample manipulation, compared to conventional infrared methods.

Additional photochemical results are consistent with the dissociative loss of CO from the surface. First, irradiation of the (S)—SiCo(CO)<sub>4</sub> under vacuum leads to disappearance of all metal carbonyl species. This disappearance is dramatically suppressed under CO, as expected. Under 1 atm Ar, irradiation leads to some decomposition, but the rate eventually declines, presumably due to the accumulation of CO in the cell. When decomposition is detected, CO and  ${\rm CO}_{2}$  are observed as gas phase photoproducts. Irradiation of the (S)-SiCo $(CO)_4$  under 1 atm  $O_2$  leads to more rapid decline of the metal carbonyl signals, and a larger signal for gas phase  ${\rm CO_2}$  is observed, compared to reaction under Ar, Figure 2. Thus,  $\boldsymbol{0}_2$  is a reactive gas and presumably interacts with the photogenerated, coordinatively unsaturated  $-\hat{S}iCo(CO)_n$  species to lead to the oxidation of CO to CO<sub>2</sub>. We estimate the  $CO/CO_2$  product ratio to be ~3/1 when the irradiation is carried out under 1 atm  $0_2$ . Irradiation of  $S = SiCo(CO)_4$  under  $^{18}O_2$  leads to the formation of  $c^{18}0^{16}0$  and irradiation of  $c^{13}$ CO-enriched  $s^{16}$ CO<sub>2</sub> under  $c^{18}$ O<sub>2</sub> leads to  $^{13}$ C $^{18}$ O $^{16}$ O. These results establish the source of CO $_2$  to be CO originally bound to Co and  $0_2$  in the gas phase. The Co product is likely some form of Co-oxide; PAS in the visible region is consistent with this conclusion. $^{9}$ 

FTIR/PAS can also be used to monitor photosubstitution of the  $S-SiCo(CO)_4$  using gas phase entering groups such as  $PF_3$  and olefins, Figure 3. At 25°C, the chemistry represented by (5) is detectable. The main

 $S = SiCo(CO)_4 + PF_3 \xrightarrow{hv} S = SiCo(CO)_n (PF_3)_{4-n}$  (5) infrared feature is at ~2000 cm<sup>-1</sup> and is associated with  $S = SiCo(CO)_3 PF_3$  with the PF<sub>3</sub> occupying an axial position trans to the Si=0, as found for R<sub>3</sub>SiCo-(CO)<sub>3</sub>P(OPh)<sub>3</sub> prepared by photosubstitution. 1,10 Irradiation of  $S = SiCo(CO)_n (PF_3)_{4-n}$  under 1 atm CO leads to the regeneration of a large fraction of the  $S = SiCo(CO)_4$ . Irradiation of  $S = SiCo(CO)_4$  under ethylene or propylene at 25°C yields loss of all metal carbonyls; Et<sub>3</sub>SiCo(CO)<sub>3</sub>(alkene) was shown previously to decompose at 25°C. But irradiation of the

S-SiCo(CO)<sub>4</sub> at -50°C under ethylene, propylene, isobutylene, or propyne and monitored by FTIR/PAS at that temperature, leads to the generation of new surface metal carbonyl species, Figure 3. The signal at ~1990 cm<sup>-1</sup> would logically be attributed to S-SiCo(CO)<sub>3</sub>(alkene), where the alkene is trans to the -Si-, paralleling solution studies. A signal that grows in at ~2050 cm<sup>-1</sup> for ethylene, propylene, or propyne, but not isobutylene, is unidentified at this time. Such a signal could be due to another isomer of the S-SiCo(CO)<sub>3</sub>(alkene) where the alkene is cis to the -Si-, or the peak could be due to multiple substitution. This point is currently under investigation, but it does appear that the species associated with the ~2050 cm<sup>-1</sup> peak is a primary product, since it appears even at short irradiation times. Interestingly, the ~2050 cm<sup>-1</sup> peak does not appear in low temperature irradiations of Ph<sub>3</sub>SiCo(CO)<sub>4</sub> in solutions containing 1-pentene or ethylene, but the Ph<sub>3</sub>Si- group may offer more steric restriction than the S-Si-.

To summarize, we have established the primary photoreaction of  $S \to SiCo(CO)_4$  to be loss of CO. This can be exploited to study the oxidation of CO by  $O_2$ , to prepare inert or labile substitution products, and to elaborate the mechanism of the reactions of immobilized, photogenerated, coordinatively unsaturated intermediates. The technique of FTIR/PAS is shown to be a molecular specific tool having unique capabilities for the study of photoreactions of the surface species on high surface area supports.

Acknowledgements. MSW and CLR acknowledge the support of the Office of Naval Research and the Dow Chemical Company and RHS and JBK acknowledge the support of the National Science Foundation, Grant No. DMR 78-24185. We thank the National Institutes of Health, Grant No. GM 27551, for support for the Nicolet 7199 FTIR used in these studies.

### References

- (a) Reichel, C.L.; Wrighton, M.S. <u>Inorg. Chem.</u>, <u>1980</u>, <u>19</u>, 3385;
   (b) Reichel, C.L.; Wrighton, M.S. <u>J. Am. Chem. Soc.</u>, submitted for publication.
- 2. (a) Sanner, R.D.; Austin, R.G.; Wrighton, M.S.; Honnick, W.D.; Pittman, Jr., C.U. Adv. Chem. Ser., 1978, 184, 13; Inorg. Chem., 1979, 18, 928; (b) Hucul, D.A.; Arnold, T.; Brenner, A. 179th ACS Meeting, INOR 7, Houston, Texas, 1980.
- 3. (a) Rockley, M.G. Chem. Phys. Lett., 1979, 68, 455; 1980, 75, 370; Appl. Spec., 1980, 34, 405, 407; (b) Rockley, M.G.; Davis, D.M.; Richardson, H.H. Science, 1980, 210, 918; (c) Vidrine, D.W. Appl. Spec., 1980, 34, 314; (d) Royce, B.S.H.; Enus, J.; Teng, Y.C. Bull. Am. Phys. Soc., 1980, 25, 408; (e) Laufer, G.; Juneke, J.T.; Royce, B.S.H.; Teng, Y.C. Appl. Phys. Lett., 1980, 37, 517; (f) Royce, B.S.H.; Teng, Y.C.; Enus, J. IEEE Symp. on Ultrasonics, 1980, Vol. 2, 652.
- 4. There are a number of techniques useful for obtaining vibrational spectra of surface species, cf. "Vibrational Spectroscopies for Adsorbed Species", A. T. Bell, M.L. Hair, Eds., <u>ACS Symposium Series</u>, 1980, 137, American Chemical Society, Washington, D.C. The FTIR/PAS would appear to offer advantages when monitoring both surface and gas phase reaction products in <u>situ</u>.
- In a typical procedure, ~1 g of high surface area  ${\rm SiO}_2$  having some -OH available is suspended in ~25 ml of alkane solvent containing excess (EtO)\_3SiH at 25°C under N\_2. The solution is stirred for 24 h and then filtered to collect a solid. After repeated washing the solid exhibits an infrared signal at ~2250 cm<sup>-1</sup> associated with Si-H. Infrared transmission experiments indicate that there is ~6.1 mol per cent  $-{\rm SiH}$ . The powder is then reacted with excess  ${\rm Co}_2({\rm CO})_8$  in alkane solution under N<sub>2</sub> at 25°C for 24 h. After repeated washing, the solid still exhibits an infrared peak at ~2250 cm<sup>-1</sup> (~3.4 mol per cent SiH) and signals at 2110, 2050, and ~2020 cm<sup>-1</sup> characteristic of  $-{\rm SiCo}({\rm CO})_4$  (~1.9 mol percent). The Reaction of the powder with  ${\rm Co}_2({\rm CO})_8$  also leads to some decomposition of the  ${\rm Co}_2({\rm CO})_8$  to leave a Co-oxide/hydroxide on the surface. The Samples were exposed to vacuum for 20 min at 25°C prior to any experimentation to remove any 0<sub>2</sub> or H<sub>2</sub>0 from the surface.
- 6. (a) Arkles, B. <u>Chemtech</u>, <u>1977</u>, <u>7</u>, 766; (b) Whitehurst, D.D. <u>Chemtech</u>, <u>1980</u>, <u>10</u>, 44; (c) Karger, B.L.; Giese, R.W. <u>Anal. Chem.</u>, <u>1978</u>, <u>50</u>, 1048A; (d) Grushka, E.; Kikta, E.J. <u>Anal. Chem.</u>, <u>1977</u>, <u>49</u>, 1004A; (e) Murray, R.W. <u>Accs. Chem. Res.</u>, <u>1980</u>, <u>13</u>, 135.
- 7. Chalk, A.J.; Harrod, J.F. J. Am. Chem. Soc., 1967, 39, 1640.


- 8. FIIR/PAS spectra were obtained using a Nicolet 7199 FIIR with a gas microphone photo-acoustic detector of our own design that allows control of the sample environment. The details are to be published independently by JBK and RHS. Data collection for 8 cm<sup>-1</sup> resolution spectra typically take 4 min. The acoustic coupling gas in all cases was 1 atm Ar (<1 ppm H<sub>2</sub>O, O<sub>2</sub>) except when monitoring gas phase products as in photolyses under O<sub>2</sub>.
- 9. The Co-oxide product is detected as an absorber in the visible at ~620 nm. <sup>1b</sup> Monitoring a sample by FTIR/PAS and by PAS in the visible shows the decline of metal carbonyl to be accompanied by the growth of the ~620 nm feature attributed to Co-oxide.
- 10. Many metal carbonyls undergo photosubstitution under PF3, Geoffroy, G.L.; Wrighton, M.S. "Organometallic Photochemistry", Academic Press: New York, 1979, including Co carbonyls related to those under study here: Udovich, C.A.; Clark, R.J. <u>Inorg. Chem.</u>, 1969, 8, 938.


### Figure Captions


Figure 1. FTIR/PAS spectrum before, curve 1, and after, curve 2, 20 min near-uv irradiation of S— $SiCd(CO)_4$  under 1 atm  $^{13}CO$ . In each case the acoustic coupling gas is 1 atm Ar. The spectral changes are consistent with the photochemical generation of S— $SiCo(^{13}CO)_4$ . No  $^{13}CO$  exchange occurs in the dark at  $25^{\circ}C$  on the timescale of this experimentation.

Figure 2. FTIR/PAS spectrum before, curve 1, and after, curve 2, 20 min near-uv irradiation of S  $SiCo(CO)_4$  under 1 atm  $O_2$ . For spectra 1 and 2 the acoustic coupling gas is the 1 atm  $O_2$ . Spectrum 2 shows gaseous CO and  $CO_2$  photoproducts. Spectrum 3 is after evacuating the sample chamber to remove all gases and introducing  $O_2$  at 1 atm as the acoustic coupling gas.

Figure 3. (A) Curve 1 is the initial FTIR/PAS spectrum of a sample of  $S-SiCo(CO)_4$  and curve 2 is the spectrum obtained after near-uv irradiation, of the sample characterized in 1, for 10 min under 1 atm PF<sub>3</sub>. The product is  $S-SiCo(CO)_n(PF_3)_{4-n}$ . Curve 3 is the spectrum that results when the sample characterized in 2 is irradiated under 1 atm of CO to regenerate  $S-SiCo(CO)_4$ . These spectra are recorded using 1 atm Ar as the coupling gas. (B) Spectral changes accompanying the near-uv irradiation of  $S-SiCo(CO)_4$ , curve 1, at low temperature,  $\sim -50^{\circ}C$ , under  $\sim 100$  torr isobutylene for 10 min to form  $S-SiCo(CO)_3$  (isobutylene), curve 2.







### TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies		No. Copies
Office of Naval Research		U.S. Army Research Office	
Attn: Code 472		Attn: CRD-AA-IP	
800 North Quincy Street		P.O. Box 1211	
Arlington, Virginia 22217	2	Research Triangle Park, N.C. 27709	1
ONR Branch Office		Naval Ocean Systems Center	
Attn: Dr. George Sandoz		Attn: Mr. Joe McCartney	
536 S. Clark Street		San Diego, California 92152	I
Chicago, Illinois 60605	1	•	
•		Naval Weapons Center	
OM trea Offige		Attn: Dr. A. B. Amster,	
AT TO A PROPERTY COME.		Chemistry Division	
713 Broadway		China Lake, California 93555	1
New York, New York 10005	1	·	
		Naval Civil Engineering Laboratory	
ONR Western Regional Office		Attn: Dr. R. W. Drisko	
1030 East Green Street		Port Hueneme, California 93401	1
Pasadena, California 91106	1		
		Department of Physics & Chemistry	
ONR Eastern/Central Regional Office		Naval Postgraduate School	
Attn: Dr. L. H. Peebles		Monterey, California 93940	1
Building 114, Section D			
666 Summer Street	_	Dr. A. L. Slafkosky	
Boston, Massachusetts 02210	1	Scientific Advisor	
		Commandant of the Marine Corps	
Director, Naval Research Laboratory		(Code RD-1)	1
Attn: Code 6100	-	Washington, D.C. 20380	1
Washington, D.C. 20390	1	Office of Newal Because	
The Indones Company		Office of Naval Research Attn: Dr. Richard S. Miller	
The Assistant Secretary			
of the Navy (RE&S)		800 N. Quincy Street Arlington, Virginia 22217	1
Department of the Navy Room 4E736, Pentagon		Allington, virginia 22217	•
Washington, D.C. 20350	1	Naval Ship Research and Development	
washington, Dece 20000	•	Center	
Commander, Naval Air Systems Command	i	Attn: Dr. G. Bosmajian, Applied	
Attn: Code 310C (H. Rosenwasser)	•	Chemistry Division	
Department of the Navy		Annapolis, Maryland 21401	1
Washington, D.C. 20360	1		
		Naval Ocean Systems Center	
Defense Technical Information Center	•	Attn: Dr. S. Yamamoto, Marine	
Building 5, Cameron Station		Sciences Division	
Alexandria, Virginia 22314	12	San Diego, California 91232	1
Dr. Fred Saalfeld		Mr. John Boyle	
Chemistry Division, Code 6100		Materials Branch	
Naval Research Laboratory		Naval Ship Engineering Center	
Washington, D.C. 20375	1	Philadelphia, Pennsylvania 19112	1

# TECHNICAL REPORT DISTRIBUTION LIST, GEN

No.	
Copie	S

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503 1
Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402 1

# TECHNICAL REPORT DISTRIBUTION LIST, 359

	No. Copies		No. Copies
Dr. Paul Delahav		Dr. P. J. Hendra	
Department of Chemistry		Department of Chemistry	
New York University		University of Southhampton	
New York, New York 10003	1	Southhampton SO9 5NH	
		United Kingdom	1
Dr. E. Yeager			
Department of Chemistry		Dr. Sam Perone	
Case Western Reserve University		Department of Chemistry	
Cleveland, Ohio 41106	1	Purdue University	
		West Lafayette, Indiana 47907	1
Dr. D. N. Bennion			
Department of Chemical Engineering		Dr. Royce W. Murray	
Brigham Young University		Department of Chemistry	
Provo, Utah 84602	1	University of North Carolina	
		Chapel Hill, North Carolina 27514	1
Dr. R. A. Marcus			
Department of Chemistry		Naval Ocean Systems Center	
California Institute of Technology	•	Attn: Technical Library	_
Pasadena, California 91125	1	San Diego, California 92152	1
Dr. J. J. Auborn		Dr. C. E. Mueller	
Rell Laboratories		The Electrochemistry Branch	
Murray Hill, New Jersey 07974	1	Materials Division, Research & Technology Department	
Dr. Adam Heller		Naval Surface Weapons Center	
Bell Laboratories		White Oak Laboratory	
Murray Hill, New Jersey 07974	1	Silver Spring, Maryland 20910	1
,,,,,,	•	officer opting, haryland 20010	•
Dr. T. Katan		Dr. G. Goodman	
Lockheed Missiles & Space		Globe-Union Incorporated	
Co, Inc.		5757 North Green Bay Avenue	
P.O. Box 504		Milwaukee, Wisconsin 53201	1
Sunnyvale, California 94088	1		
		Dr. J. Boechler	
Dr. Joseph Singer, Code 302-1		Electrochimica Corporation	
NASA-Lewis		Attention: Technical Library	
21000 Brookpark Road	•	2485 Charleston Road	_
Cleveland, Ohio 44135	1	Mountain View, California 94040	1
Dr. B. Brummer		Dr. P. P. Schmidt	
EIC Incorporated		Department of Chemistry	
55 Chapel Street		Oakland University	
Newton, Massachusetts 02158	1	Rochester, Michigan 48063	1
Library		Dr. H. Richtol	
P. R. Mallory and Company, Inc.		Chemistry Department	
Northwest Industrial Park		Rensselaer Polytechnic Institute	
Burlington, Massachusetts 01803	1	Troy, New York 12181	1

# TECHNICAL REPORT DISTRIBUTION LIST, 359

	No.			No. Copies
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1	Dr. R. P. Van Duyne Department of Chemistry Northwestern University Evanston, Illinois 60201	l	1
Pr. M. Wrighton Chemistry Department Massach Setts Institute of Technology Cambridge, Massachusetts 02139	ì	Dr. B. Stanley Pons Department of Chemistry University of Alberta Edmonton, Alberta CANADA T6G 2G2	1	1
Larry E. Plew Naval Weapons Support Center Code 30736, Building 2906 Crane, Indiana 47522	1	Dr. Michael J. Weaver Department of Chemistry Michigan State University East Lansing, Michigan 48824		1
S. Rubv DOE (STOR) 600 E Street Washington, D.C. 20545	1	Dr. R. David Rauh EIC Corporation 55 Chapel Street Newton, Massachusetts 02158		1
Dr. Aaron Wold Brown University Department of Chemistry Providence, Rhode Island 02192	I	Dr. J. David Margerum Research Laboratories Division Hughes Aircraft Company 3011 Malibu Canyon Road Malibu, California 90265		1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1	Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton 509 5NH England		1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1	Dr. Janet Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214		1
Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California	1	Dr. R. A. Ostervoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214		1
Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratorv Silver Spring, Marvland 20910	1	Mr. James R. Moden Naval Underwater Systems Center Code 3632		-
		Newport, Rhode Island 02840		1

## TECHNICAL REPORT DISTRIBUTION LIST, 359

	No.		No.
	Copies		Copies
Dr. R. Nowak		Dr. John Kincaid	1
Naval Research Laboratory		Department of the Navy	
Code 6130		Stategic Systems Project Office	
Washington, D.C. 20375	1	Room 901	
		Washington, DC 20376	
Dr. John F. Houlihan		•	
Shenango Valley Campus		M. L. Robertson	
Pennsylvania State University		Manager, Electrochemical	
Sharon, Pennsylvania 16146	1	Power Sonices Division	
		Naval Weapons Support Center	
Dr. M. G. Sceats		Crane, Indiana 47522	1
Department of Chemistry		•	
University of Rochester		Dr. Elton Cairns	
Rochester, New York 14627	1	Energy & Environment Division	
		Lawrence Berkeley Laboratory	
Dr. D. F. Shriver		University of California	
Department of Chemistry		Berkeley, California 94720	1
Northwestern University		• •	
Evanston, Illinois 60201	1	Dr. Bernard Spielvogel	
		U.S. Army Research Office	
Dr. D. H. Whitmore		P.O. Box 12211	
Department of Materials Science		Research Triangle Park, NC 27709	1
Northwestern University		·	
Evanston, Illinois 60201	1	Dr. Denton Elliott	
		Air Force Office of	
Dr. Alan Bewick		Scientific Research	
Department of Chemistry		Bldg. 104	
The University		Bolling AFB	
Southampton, SO9 5NH England	1	Washington, DC 20332	1
Dr. A. Himy			
NAVSEA-5433	•		
NC #4			
2541 Jefferson Davis Highway			
Arlington, Virginia 20362	1		

# TECHNICAL REPORT DISTRIBUTION LIST, 051A

	No. Copies		No. Copies
Dr. M. A. El-Sayed		Dr. M. Rauhut	
Department of Chemistry		Chemical Research Division	
University of California,		American Cyanamid Company	
Los Angeles		Bound Brook, New Jersey 08805	1
Los Angeles, California 90024	1		
		Dr. J. I. Zink	
Dr. E. R. Bernstein		Department of Chemistry	
Department of Chemistry		University of California,	
Colorado State University		Los Angeles	_
Fort Collins, Colorado 80521	1	Los Angeles, California 90024	1
Dr. C. A. Heller		Dr. D. Haarer	
Naval Weapons Center		IBM	
Code 6059		San Jose Research Center	
China Lake, California 93555	1	5600 Cottle Road	
		San Jose, California 95143	1
Dr. J. R. MacDonald			
Chemistry Division		Dr. John Cooper	
Naval Research Laboratory		Code 6130	
Code 6110		Naval Research Laboratory	
Washington, D.C. 20375	1	Washington, D.C. 20375	1
Dr. G. B. Schuster		Dr. William M. Jackson	
Chemistry Department		Department of Chemistry	
University of Illinois		Howard University	
Urbana, Illinois 61801	1	Washington, DC 20059	1
Dr. A. Adamson		Dr. George E. Walraffen	
Department of Chemistry		Department of Chemistry	
University of Southern		Howard University	
California		Washington, DC 20059	1
Los Angeles, California 90007	1		
Or. M. S. Wrighton			
Department of Chemistry			
Massachuse Institute of			
Technology			
Carbridge, Massachusetts 02139	1		

# DATE